

EXHIBIT B

This data is from an experiment examining the ability of anti-PDGF IgG to neutralize the chemotactic activity present in the Anti-PDGF IgG purified fraction (PDGF related) of human wound fluids collected on various days (D1 = Day 1, D2 = Day 2, etc.) As seen in the column to the far right only day 1 shows significantly high activity which is neutralizable with the anti-PDGF IgG. ie sample 1 is Day 1 in the presence of non-immune IgG 157; sample 2 is the same material in the presence of anti-PDGF IgG 28. Sample 19 is a 10 ng/ml PDGF standard in the presence of non-immune IgG 167; sample 20 is the same material in the presence of anti-PDGF IgG 40. Therefore samples below with activity below 28-40 have little or no PDGF like activity.

				ABS	Activity, μ m
1	D1	IgG	31	249	157
2	"	α PDGF	61	124	28
3	D2	IgG	33	125	29
4		α PDGF	17	108	12
5	D3	IgG	10	126	30
6		α PDGF	89	76	-
7	D4	G	1	86	-
8		P	60	45	-
9	D5	G	66	99	3
10		P	9	48	-
11	D6	G	11	105	9
12		P	74	76	-
13	D7	G	35	108	22
14		P	21	110	14
15	D8	B	97	123	27
16		P	68	107	11
17	D9	G	44	103	7
18		P	93	93	-
19	PDGF 10ng/ml	IgG	81	265	167
20		α PDGF	84	136	40
21	5ng/ml	IgG	15	197	101
22		α PDGF	78	104	8
23		P	28	92	-